

UNCLASSIFIED


LA-1325

Subject Category: PHYSICS

UNITED STATES ATOMIC ENERGY COMMISSION

THE CRYSTAL STRUCTURE OF  
GAMMA PLUTONIUM


By  
W. H. Zachariasen



September 1951

Los Alamos Scientific Laboratory  
Los Alamos, New Mexico

Technical Information Extension, Oak Ridge, Tennessee



19950815 066

UNCLASSIFIED

DTIC QUALITY INSPECTED 8

Date Declassified: March 15, 1956.

#### LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, express or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission to the extent that such employee or contractor prepares, handles or distributes, or provides access to, any information pursuant to his employment or contract with the Commission.

This report has been reproduced directly from the best available copy.

Issuance of this document does not constitute authority for declassification of classified material of the same or similar content and title by the same authors.

Printed in USA, Price 15 cents. Available from the Office of Technical Services, Department of Commerce, Washington 25, D. C.

LOS ALAMOS SCIENTIFIC LABORATORY  
of the  
UNIVERSITY OF CALIFORNIA

Report Written:  
September 1951

LA-1325

THE CRYSTAL STRUCTURE  
OF GAMMA PLUTONIUM

by  
W. H. Zachariasen

Work performed under Contract No. W-7405-Eng-36

Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

# ABSTRACT

Gamma plutonium is found to be orthorhombic with eight atoms in a unit cell of dimensions (at 226°C)  $a_1 = 3.1518$  kX,  $a_2 = 5.7568$  kX,  $a_3 = 10.142$  kX. The calculated density is 17.13. The space group is Fddd and the positions of the eight atoms are: (0 0 0) (0 1/2 1/2) (1/2 0 1/2) (1/2 1/2 0) (1/4 1/4 1/4) (1/4 3/4 3/4) (3/4 1/4 3/4) (3/4 3/4 1/4).

Each plutonium atom is bonded to ten others at an average distance of 3.152 kX, four being at 3.020 kX, two at 3.152 kX and four at 3.285 kX.

The coefficients of thermal expansion are found to be  $\alpha_{100} = -22.5 \times 10^{-6}$ ,  $\alpha_{010} = +47 \times 10^{-6}$ ,  $\alpha_{001} = +94 \times 10^{-6}$

## THE CRYSTAL STRUCTURE OF GAMMA PLUTONIUM

By

W. H. Zachariasen

The crystal structure is known for only two of the five allotropic forms of plutonium metal. The delta modification is cubic face-centered<sup>1</sup> with  $a = 4.63 \text{ \AA}$ <sup>2</sup> (at 350°C), and the epsilon form is cubic body-centered with  $a = 3.63 \text{ \AA}$ <sup>3</sup> (at 510°C).

Satisfactory X-ray diffraction patterns of the alpha, beta and gamma forms of plutonium have been available for several years. However, the indexing of these powder patterns has proved to be a very difficult task. The indexing of the pattern of gamma plutonium given in this report succeeded only after a great many hours of intensive work. The patterns of alpha and beta plutonium have not yet been interpreted.

Table 1 gives the observed intensities and sine squares as obtained from a powder diffraction pattern of gamma plutonium taken with  $\text{CuK}\alpha$ -radiation at 226°C. The  $\alpha_1\alpha_2$ -doublet is separated for  $\sin^2\theta > 0.400$ , and the measurements refer to the stronger component.

The observed sine squares fit the quadratic form

$$\sin^2\theta = 0.05948 H_1^2 + 0.01728 H_2^2 + 0.005745 H_3^2.$$

The corresponding orthorhombic unit cell has dimensions  $a_1 = 3.1518 \pm 0.0004 \text{ kX}$ ,  $a_2 = 5.7568 \pm 0.0008 \text{ kX}$ ,  $a_3 = 10.142 \pm 0.002 \text{ kX}$ . Since the experimentally determined density is 17.0, the unit cell contains eight atoms. The calculated density is  $\rho = 17.13$ .

The translation lattice is face-centered as shown by the fact that reflections are present only from planes for which  $H_1 H_2 H_3$  are either all even or all odd. It is further seen from Table 1 that reflections are missing from planes with  $H_1 H_2 H_3$  all even unless  $\sum H_j = 4n$ .

According to these systematic absences the plutonium atoms are arranged on two interpenetrating face-centered lattices displaced relative to each other by one-fourth of the body diagonal. The positions of the eight atoms in the unit cell are thus: (0 0 0) (0 1/2 1/2)

TABLE 1. X-RAY DIFFRACTION DATA FOR GAMMA PLUTONIUM (226°C).

Int. obs.	$\sin^2 \theta$		$H_1H_2H_3$	Int.		$H_1H_2H_3$	$\sin^2 \theta$		Int. calc.
	obs.	calc.		obs.	calc.		obs.	calc.	
s	0.0836	0.0830	111	142	w-*	155	6490	6489	8
w	925	919	004	61	w*	048	6526	6530	8
m+	949	943	022	115	w-	062	6648	6649	8
m	1297	1290	113	80	w+	228	6766	6769	15
m-	2218	2209	115	36	vw+	139	6853	6853	8
w+	2265	2257	131	36	vw+	315	6970	6967	8
w+	2616	2609	202	28	w*	331	7011	7015	8
w+	2723	2717	133	26	m-	246	7300	7300	15
w+	2788	2781	026	25	w-	333	7473	7475	8
vw	2860	2853	040	12	vw	1.1.11	7724	7724	8
w+	3100	3092	220	21	w-	157	7870	7868	8
w	3590	3588	117	17	w-	2.0.10	8125	8124	8
w	3637	3636	135	16	vw-	0.0.12	8273	8273	4
vw	3687	3677	008	8	w-	317	8345	8346	9
w+	3772	3772	044	15	w+*	335	8392	8394	9
m	4012	4011	224	30	w	066	8486	8487	9
w	4448	4447	206	12	w	260	8800	8798	10
w-	5017	5015	137	9	w-	1.3.11	9150	9151	12
w-	5111	5110	151	9	w	171	9388	9390	13
vw	5427	5426	119	9	vw	400	9517	9517	8
m*	5461	5462	242	18	ms	159	9717	9706	19
vw	5574	5570	153	9	m*	264	9768	9717	38
w*	5597	5588	311	9	w+	337	9849	9773	21
vw+	6046	6048	313	8	w+	173	9850	9850	27
vw+	6458	6458	0.2.10	8	w+	351	9867	9868	29

\* Coincidence with  $\alpha_2$  - line.

(1/2 0 1/2) (1/2 1/2 0) (1/4 1/4 1/4) (3/4 3/4 1/4) (3/4 1/4 3/4) (1/4 3/4 3/4). This atomic configuration corresponds to the space group symmetry Fddd ( $D_{2h}^{24}$ ) with the eight plutonium atoms structurally equivalent. A perspective drawing of one unit cell is shown in Figure 1.

The last column of Table 1 gives the intensities calculated from the formula

$$I \propto |F|^2 \frac{1 + \cos^2 2\theta}{\sin^2 \theta \cos \theta}$$

The absorption and temperature factors have not been taken into account. Observed and calculated intensities should accordingly be compared only for neighboring reflections.

The crystal structure of gamma plutonium is unlike that of any other metal. Each plutonium atom is bonded to ten others at approximately the same distance, as illustrated in Figure 2. The individual interatomic distances are (at 226°C): Pu - 4Pu = 3.020 kX, Pu - 2Pu = 3.152 kX, Pu - 4Pu = 3.285 kX. The average distance of Pu - 10Pu = 3.16Å compares to Pu - 12Pu = 3.27Å in delta plutonium and Pu - 8Pu = 3.15Å in epsilon plutonium.

When allowance is made for the effect of coordination number one finds a metallic radius of 1.60Å for plutonium in the gamma form as compared to the radius 1.63Å in delta plutonium. The difference in radius for the two forms is possibly due to the fact that the transition from the delta to the gamma form is accompanied by the promotion of a fraction of an electron from the 5f to the 6d level. From the value of 1.60Å for the radius of plutonium in the gamma form, one would estimate that there are about three electrons in the 5f shell.

Normal to the  $a_3$ -axis there are pseudo-hexagonal layers, the  $a_3$ -period corresponding to four times the layer separation. The bond lengths within a layer are 3.152 kX and 3.285 kX. The bond length between layers is 3.020 kX.

Table 2 shows the variation of the unit cell dimensions with temperature in the range 161°C - 265°C, and Table 3 gives the deduced values of the three principal coefficients of thermal expansion.




Fig. 1. The Unit Cell of Gamma Plutonium.






Fig. 2. A portion of the unit cell illustrating the tenfold coordination. Relative to Figure 1, the origin of the unit cell has been displaced by  $(0\ 0\ 1/2)$ .

TABLE 2. UNIT CELL DIMENSIONS (IN kX)  
AND CALCULATED DENSITY AT  
VARIOUS TEMPERATURES

	161°	200°	226°	265°	Accuracy
$a_1$	3.1568	3.1527	3.1518	3.1490	$\pm 0.0004$ kX
$a_2$	5.7400	5.7475	5.7568	5.7670	$\pm 0.0008$
$a_3$	10.083	10.111	10.142	10.180	$\pm 0.002$
$\rho$	17.26	17.21	17.13	17.06	

TABLE 3. COEFFICIENTS OF THERMAL EXPANSION

$10^6 \alpha_{100}$	$- 22.5 \pm 1.0$
$10^6 \alpha_{010}$	$+ 47 \pm 2$
$10^6 \alpha_{001}$	$+ 94 \pm 4$

## REFERENCES

- <sup>1</sup> R. C. L. Mooney and W. H. Zachariasen, CK-1367, CK-1377, CK-1968 and CN-2069.
- <sup>2</sup> F. J. Schnettler and E. R. Jette, LAMS-211, p. 21. F. H. Ellinger of Los Alamos Scientific Laboratory has obtained evidence that  $4.624 \pm 0.001$  kX is a somewhat better value for the unit cell edge of delta plutonium at  $350^{\circ}\text{C}$ . However, there is good reason to suspect that the specific volume of the delta phase at any given temperature is strongly dependent on previous history, and it is not clear that "equilibrium" values of unit cell dimensions have ever been determined.
- <sup>3</sup> F. J. Schnettler and E. R. Jette, LAMS-249, p. 17.

## ACKNOWLEDGEMENT

The X-ray diffraction patterns upon which this investigation is based were taken by Finlay Ellinger and measured by Marian Gibbs.

The author is also indebted to Eric R. Jette, CMR Division, and to A. S. Coffinberry, Group CMR-5, for having provided the opportunity for this investigation to be carried out.